



Research paper

Acute Toxicity of Potassium Sulphate in Amami Rabbit (*Pentalagus furnessi*) with Reference to Biochemical Attributes

Ibiama Kenneth Adonye ^a, Solaja Mayowa ^a, Adeola Ogunrin ^{b,*}

^a Department of Biological Sciences, Federal University of Technology, PMB 1526, Owerri, Nigeria

^b Department of Medicine, Clinical School, International Medical University, Jalan Rasah, 70300 Seremban, Malaysia

ARTICLE INFO	ABSTRACT
<p><i>Article history</i></p> <p>Received 16 November 2020 Revised 18 October 2021 Accepted 22 November 2021 Published 02 December 2021</p> <hr/> <p><i>Keywords</i></p> <p>Serum Biochemical alteration Potassium sulfate Rabbit <i>Pentalagus</i></p>	<p>The aim of present study was to see how acute potassium sulphate toxicity affected Amami rabbit biochemical parameters. Protein and cholesterol levels in rabbits treated with potassium sulphate decreased, while creatinine and bilirubin levels increased. Protein (-5.5 to -20.0 percent), cholesterol (-6.7 to -13.3 percent), creatinine (8.3 to 16.6 percent), and bilirubin (2.2 to 6.7 percent) were also measured during the exposure period. The findings point to a change in blood serum biochemicals as a result of potassium sulphate exposure. These metrics are important indications of a live organism's health and stress.</p>

1. Introduction

Both developed and developing countries that are progressing rapidly in the field of agriculture, biotechnology and industry introducing various kinds of harmful substances in to the environment and thereby facing a serious problem. In the agriculture industries artificial ripeners are used to ripener fruits and vegetables in the processes as

transport, packaging, storage, pre- and post –harvest ripening, etc. Nowadays artificial ripeners such as calcium carbide, acetylene gas, carbon monoxide, potassium sulfate, zyme, ethephon, potassium dihydrogen arthoposphate, etc. are used for fruit ripening especially on fruits including apricots, mango, bananas, papayas, plums, cherry,

* Corresponding author: Adeola Ogunrin, E-mail: o_adeola@gmail.com



DOI

<https://doi.org/10.5281/zenodo.5807840>



pomegranate, Japanese fruits, etc. These artificial ripener known as silent killer because of their impact on memory loss, cerebral edema, colonic, prostates and lung cancer, quick-buck syndrome, DNA, RNA and hematological changes. Potassium sulfate (K_2SO_4) is being used as pre-harvest ripener on mango, pineapple, coffee, tomato, cucumber, groundnut and rubber. It reported toxicity, flammability, destructive, self-reactive and incompatible with water oxidizing and other symptoms (Malik et al., 2021). Due to the adverse effect of this artificial ripener present study was assigned on Amami rabbit, *Pentalagus furnessi*.

2. Materials and Methods

2.1 Experimental Animal

Normal adult healthy Amami rabbit, *Pentalagus furnessi* weighting from 2.25 ± 0.12 kg was used in the experiment. The animals were purchased from Reena Rabbitry, Rahata and acclimatized at room temperature ($26 \pm 20^\circ C$). Experimental protocol was approved by the Institutional Animal Ethics Committee (IAEC). They were maintained (55 cm L x 45 cm W x 30 cm H) in cage with a constant 12 hours light and dark rhythms. The control group provided water and food while treatment rabbit provided a food with acute dose at 5 mg/kg body weight (Ramalingam et al., 2001). The post treatment followed up to 28 day.

2.2 Experimental Treatment

Potassium sulfate was dissolved in 2 ml distilled water. An acute oral dose was performed according to the Office of Prevention, Pesticide and Toxic Substances (OPPTS) guidelines following the limit test procedure. Animals were fasted overnight prior to working.

2.3 Biochemical Analysis

Blood sample was collected from ear artery using heparinized syringes (5.000 UI) in sterilized vial on control 7, 14, 21 and 28 days. The following biochemical parameters were estimated according to standard method as: protein (Lowry et al., 1951), cholesterol (Zlatkis et al., 1953), bilirubin (Osar, 1965) and creatinine (Kokka & George, 1970).

3. Results and Discussion

An observation made on blood serum biochemical contents indicates that the protein and cholesterol get decline while creatinine and bilirubin incline after the intoxication with potassium sulfate (Figure 1- 4).

3.1 Protein

Liver is responsible for the synthesis of plasma protein. After treated it showed minimum (-5.5%) and maximum (-20.0%) decline after intoxication on 28 and 7 days (Figure 1). The general trend of decline was 7th toward 28th days. Reduction in serum total protein in the present study may ascribe to lesser food intake, poor nutrition and liver malfunction of treated animal. Similar decline in total protein have been reported (Sharma et al., 2007) after administration of textile dye waste water to rats. Several opinions have been put forward to explain it as decline in protein level showed the tress condition of the animal (Rajmannar & Manohar, 1998). The depletion in protein may be due to proteolytic (Blum et al., 1985; Horai et al., 2006). Toxicant influences the change in tissue protein into soluble functions lead to blood for use (Rao et al., 1990). The reduction in protein is further comparable, increased proteolysis, shifting in nitrogen metabolism (Ramalingam, 1982). Inhibition of ribosomal activity resulted in protein degradation are also being the possible reasons for depletion. Reduction in protein might be due to the metabolic stress caused by toxicant leads to depletion in of protein synthesis (Shakoori et al., 1976). Under stress condition it supplied energy to metabolic pathway and biochemical reactions (Rekha & Kumar, 2008). Another reason for depletion might be due to the low availability of amino acids, which would be used for fatty acid and cholesterol synthesis (Ohnishi et al., 2017; Yeragi et al., 2000).




Figure 1. Showing changes in protein

3.2 Creatinine

In the present study the level of creatinine increased significantly. The level of creatinine after two week exposure showed incline up to 16.6% on 28 day (Figure 2). Similar type of incline trends was observed in the HDL, LDL and VLDL cholesterol. We know that the parameter is determined to assess the potential effects of chemical on glomerulus filtration rate (GFR). As a substance is normally filtered by the glomerulus, an increase in serum concentration thus suggests a decrease in GFR (Eaton & Klassen, 1996; Verley, 1969). Sharma et al. (2006) reported that glomerule nephrosis and degradation of tubular epithelium in kidney of rat exposed to textile dye, along with alteration in serum biochemical parameters viz. creatinine, protein, cholesterol. Thus in the present investigation the renal toxicity might have increased the serum creatinine level. Creatinine is a spontaneously formed cyclic derivative of creatinine and closely filtered out by the kidney and no reabsorption. If the filtration of the kidney is deficient, creatinine blood levels rise up. Higher levels of creatinine indicate a falling glomerular filtration rate and as a result a decreased capability of the kidneys to excrete waste products.




Figure 2. Showing changes in creatinine

3.3 Cholesterol

It has been observed that intoxication lead to reduction in cholesterol content. The depletion in cholesterol was maximum (-13.3%) and minimum in (-6.7%) at 7th and 28th day exposure to potassium sulfate respectively (Figure 3). A significant reduction in serum cholesterol, HDL, LDL, VLDL level

was found in the present study may be due to reduction in food intake and malnutrition and stress condition caused by potassium sulfate. Sharma et al. (2006) found similar result in the rat after treating with textile dye. Post treated rat did not show any recovery in serum biochemicals. Similarly Panda et al., (1987) reported a marked reduction in cholesterol level of Japanese quails and boiler chicken consequent to dietary aflatoxin. The decrease in serum cholesterol indicated degenerative changes and hypofunction of liver. The effect of intoxication to prolong the dose was found close dependent (Panda et al., 1987).




Figure 3. Showing changes in cholesterol

3.4 Bilirubin

Overall an incline was noticed in bilirubin, directly and indirectly after exposure to potassium sulfate. The level of bilirubin after two-week exposure showed rose up to maximum 6.7%, 12% and 10% in total, direct and indirect bilirubin content on 28th day respectively (Figure 4). Similar trend was also noticed in direct and indirect bilirubin contents. It is a bile pigment formed from the breakdown of haem in RBC. Bilirubin elevated level may create disease. Gilbert's syndrome and Crigler-Najjar syndrome are characterized by increased bilirubin in the serum. Hyperbilirubinemia may lead to accumulation of bilirubin in brain region leads to neurological deficits, jaundice in neonatal (Adrogué & Madias, 1981; Kubo et al., 2013), kernicterus cancer (Newman et al., 1990).




Figure 4. Showing changes in bilirubin

4. Conclusion

Present study indicates that rabbit are susceptible to potassium sulfate which produced serum biochemical alteration in blood. The blood biochemical parameters tests have been an important diagnostic tool in medical science. These parameters are valuable indicators of health and stresses in living organisms. The present work is used to establish the extent of toxicity of potassium sulfate.

Funding Information

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

Declaration of Conflict

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Research Ethics

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

References

1. Adrogué, H. J., & Madias, N. E. (1981). Changes in plasma potassium concentration during acute acid-base disturbances. *American Journal of Medicine*, 71(3), 456–467. [https://doi.org/10.1016/0002-9343\(81\)90182-0](https://doi.org/10.1016/0002-9343(81)90182-0)
2. Blum, D. C., Canaan, S., & Winkelman, J. W. (1985). *Clinical chemistry* (p. 1050) (2nd ed). Harper & Row Publishing.

3. Eaton, D. L., & Klassen, C. D. (1996). Principles of toxicology. In Klaseen (Ed.), *Toxicology. The basic science of poison*.
4. Horai, S., Minagawa, M., Ozaki, H., Watanabe, I., Takeda, Y., Yamada, K., Ando, T., Akiba, S., Abe, S., & Kuno, K. (2006). Accumulation of Hg and other heavy metals in the Javan mongoose (*Herpestes javanicus*) captured on Amamioshima Island, Japan. *Chemosphere*, 65(4), 657–665. <https://doi.org/10.1016/j.chemosphere.2006.01.078>
5. Kokka, N., & George, R. (1970). Effect of hypothalamic stimulation on blood glucose in the rabbit. *Neuroendocrinology*, 6(1), 1–9.
6. Kubo, M., Nakashima, T., Honda, T., Kochi, Y., Ito, Y., Hattori, S., & Kuraishi, T. (2013). Histopathological examination of spontaneous lesions in Amami rabbits (*Pentalagus furnessi*): A preliminary study using formalin-fixed archival specimens. *Japanese Journal of Zoo and Wildlife Medicine*, 18(2), 65–70. <https://doi.org/10.5686/jjzwm.18.65>
7. Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193(1), 265–275. [https://doi.org/10.1016/S0021-9258\(19\)52451-6](https://doi.org/10.1016/S0021-9258(19)52451-6)
8. Malik J. A., Iqbal S., Biswas J., Riaz U., & Datta S. (2021). Antidiabetic Property of *Aloe vera* (*Aloe barbadensis*) and Bitter Melon (*Momordica charantia*). In Aftab T., Hakeem K.R. (eds) *Medicinal and Aromatic Plants*. Springer, Cham. https://doi.org/10.1007/978-3-030-58975-2_10
9. Newman, T. B., Easterling, M. J., Goldman, E. S., & Stevenson, D. K. (1990). Laboratory evaluation of jaundice in newborns. Frequency, cost, and yield. *American Journal of Diseases of Children*, 144(3), 364–368. <https://doi.org/10.1001/archpedi.1990.02150270114039>
10. Ohnishi, N., Kobayashi, S., Nagata, J., & Yamada, F. (2017). The influence of invasive mongoose on the genetic structure of the endangered Amami rabbit populations. *Ecological Research*, 32(5), 735–741. <https://doi.org/10.1007/s11284-017-1489-5>
11. Osar, B. L. (1965). *Hawk's practical physiological chemistry* (14th ed) N. Y. McGraw-Hill (Ed.), 1214.
12. Panda, B. K., Praharaj, N. K., Johri, T. S., & Sah, R. L. (1987). Experimental aflatoxicosis in Japanese quails: Evidence of some biochemical changes. *Indian J. Poult. Sci*, 22, 359–362.
13. Rajamannar, K., & Manohar, L. (1998). Effect of pesticides on oxygen consumption of the fish *Labeo rohita*. *Journal of Ecobiology*, 10(3), 205–208.

14. Ramalingam, K. (1982). Effects of sublethal levels of DDT, malathion and mercury on tissue proteins of *Sarotherodon mossambicus* (Peters). *Proceedings: Animal Sciences*, 91(6), 501-505.
15. Ramalingam, V., Panneerdoss, S., Girija, M., & Ilango, S. (2001). Mercuric chloride induced changes in the histology of the testis and serum testosterone in adult albino rats. *Pollution Research*, 20(3), 439-442.
16. Rao, C. S., Chakrasali, R. T., Ila, H., & Junjappa, H. (1990). Regio-and chemoselective conjugate 1, 4-reduction of α -oxoketene dithioacetals with sodium borohydride and sodium cyanoborohydride. *Tetrahedron*, 46(6), 2195-2204.
17. Rekha, R., Gautam, R. K., Kalpana, G., & Suneel, K. (2008). Nuvan intoxication leads protein alteration in kidney of *Labeo rohita*. *Journal of Experimental Zoology, India*, 11(1), 113-115.
18. Shakoori, A. R., Zaheer, S. A., & Ahmed, M. S. (1976). Effect of malathion, dieldrin, and endrin on blood serum protein and free amino acid pool of *Channa punctatus* (bloch). *Pak. J. Zool*, 8(2), 125-134.
19. Sharma, S., Sharma, A., Singh, P. K., Soni, P., Sharma, S., Sharma, P., & Sharma, K. P. (2007). Impact of distillery soil leachate on haematology of Swiss albino mice (*Mus musculus*). *Bulletin of Environmental Contamination and Toxicology*, 79(3), 273-277. <https://doi.org/10.1007/s00128-007-9225-4>
20. Verley, H. (1969). Practical clinical biochemistry (p. 1277) (5th ed). ELBS London.
21. Yeragi, S. G., Koli, V. A., & Yeragi, S. (2000). Effect of pesticide malathion on protein metabolism of the marine crab *Uca marionis*. *Journal of Ecotoxicology & Environmental Monitoring*, 10(1), 59-62.
22. Zlatkis, A., Zak, B., & Boyle, A. J. (1953). A new method for the direct determination of serum cholesterol. *Journal of Laboratory and Clinical Medicine*, 41(3), 486-492.